

and GM and KR the Scottish data. SD undertook the main analysis and writing of the paper, with all authors, particularly JE, involved in interpretation of the results and drafting of the paper. ONG is the guarantor, who oversaw analyses at the Health Protection Agency and also commented on the drafts. SD is currently registered for a PhD at City University, London.

*Katy Sinka, Barry G Evans, Catherine M Lowndes, Neil Macdonald, Glenn Codere, David Goldberg, John V Parry and Kevin A Fenton.

REFERENCES

- 1 The UK Collaborative Group for HIV and STI Surveillance. Mapping the issues. HIV and other sexually transmitted infections in the United Kingdom in 2004. London: Health Protection Agency, Centre for Infections, 2005.
- 2 National Centre in HIV Epidemiology and Clinical Research. HIV/AIDS, viral hepatitis and sexually transmissible infections in Australia. Annual Surveillance Report 2004. National Centre in HIV Epidemiology and Clinical Research, The University of New South Wales, Sydney, NSW; Australian Institute of Health and Welfare, Canberra, ACT, 2004.
- 3 Dukers NTM, Spaargaren J, Geskus RB, *et al.* HIV incidence on the increase among homosexual men attending an Amsterdam sexually transmitted disease clinic: using a novel approach for detecting recent infections. *AIDS* 2002;**16**:F19–24.
- 4 Centers for Disease Control and Prevention. Increases in HIV diagnoses – 29 States, 1999–2002. *MMWR* 2003;**52**:1145–8.
- 5 Macdonald N, Dougan S, McGarrigle CA, *et al.* Recent trends in diagnoses of HIV and other sexually transmitted infections in England and Wales amongst men who have sex with men. *Sex Transm Infect* 2004;**80**:492–7.
- 6 Wasserheit JN. Epidemiological synergy: interrelationships between human immunodeficiency virus infection and other sexually transmitted diseases. *Sex Transm Dis* 1992;**19**:61–77.
- 7 Dodds JP, Mercey DE, Parry JV, *et al.* Increasing risk behaviour and high levels of undiagnosed HIV infection in a community sample of homosexual men. *Sex Transm Infect* 2004;**80**:236–40.
- 8 Reid D, Weatherburn P, Hickson F, *et al.* On the move: findings from the United Kingdom's gay men's sex survey 2003. London: Sigma Research, 2004.
- 9 Elford J, Bolding G, Davis M, *et al.* Trends in sexual behaviour among London homosexual men 1998–2003: implications for HIV prevention and sexual health promotion. *Sex Transm Infect* 2004;**80**:451–4.
- 10 Hart GJ, Williamson LM. Increase in HIV sexual risk behaviour in gay men in Scotland, 1996–2002: prevention failure. *Sex Transm Infect* 2005;**81**:367–72.
- 11 Mercer CH, Fenton KA, Copas AJ, *et al.* Increasing prevalence of male homosexual partnerships and practices in Britain 1990–2000: evidence from national probability surveys. *AIDS* 2004;**18**:1453–8.
- 12 Department of Health. The national strategy for sexual health and HIV. London: DOH, 2001.
- 13 Hickson F, Nutland W, Weatherburn P, *et al.* Making it count: a collaborative planning framework to reduce the incidence of HIV infection during sex between men, 3rd edn. London: Sigma Research, 2003.
- 14 Scottish Executive. Respect & responsibility—strategy and action plan for improving sexual health. Edinburgh: Scottish Executive, 2005.
- 15 Janssen RS, Satten GA, Stramer SL, *et al.* New testing strategy to detect early HIV-1 infection for use in incidence estimates and for clinical and prevention purposes. *JAMA* 1998;**280**:42–8.
- 16 Murphy G, Charlett A, Jordan LF, *et al.* HIV incidence appears constant in men who have sex with men despite widespread use of effective antiretroviral therapy. *AIDS* 2004;**18**:265–72.
- 17 Murphy G, Charlett A, Brown AE, *et al.* Is HIV incidence increasing in homo/bisexual men attending STI clinics in England, Wales and Northern Ireland? *Commun Dis Pub Health* 2004;**7**:11–14.
- 18 Dodds JP, Nardone A, Mercey DE, *et al.* Increasing high risk sexual behaviour amongst gay men in London 1996–1998: a repeated cross sectional questionnaire study. *BMJ* 2000;**320**:1510–11.
- 19 Elford J, Bolding G, Sherr L, *et al.* High-risk sexual behaviour among London gay men: no longer increasing. *AIDS* 2005;**19**:2171–4.
- 20 Mortimer JY, Salathiel JA. 'Soundex' codes of surnames provide confidentiality and accuracy in a national HIV database. *Commun Dis Rep* 1995;**15**:R183–6.
- 21 Heptonstall J, Gill ON. The legal and ethical basis for unlinked anonymous testing. *Commun Dis Rep Rev* 1989;**48**:3–6.
- 22 Medical Research Council. Guidance on collections of human tissue and biological samples for use in human research. London: Medical Research Council, 2001.
- 23 Department of Health, Clinical Ethics and Human Tissue Branch. The use of human organs and tissue—an interim statement. London: DOH, 2003.
- 24 Human tissue and biological samples for use in research: operational and ethical guidelines, 2001. <http://mrc.ac.uk> (accessed 19 Dec 2006).
- 25 PHLS Communicable Disease Surveillance Centre. Quarterly communicable disease review October to December 2001: Caldicott, Confidentiality and the Patient Information Advisory Group. *J Pub Health Med* 2002;**24**:138–44.
- 26 Dougan S, Elford J, Sinka K, *et al.* Men who have sex with men who are born abroad and diagnosed with HIV in England and Wales: an epidemiological perspective. *Int J STD AIDS* 2005;**16**:618–21.
- 27 Dougan S, Elford J, Rice B, *et al.* The epidemiology of HIV among black and minority ethnic men who have sex with men in England and Wales. *Sex Transm Infect* 2005;**81**:345–50.
- 28 Office for National Statistics. Census 2001. www.statistics.gov.uk (accessed 19 Dec 2006).
- 29 van der Bij AK, Stolte IG, Coutinho RA, *et al.* Increase of sexually transmitted infections, but not HIV, among young homosexual men in Amsterdam: are STIs still reliable markers for HIV transmission? *Sex Transm Infect* 2005;**81**:34–7.
- 30 Elford J. Changing patterns of sexual behaviour in the era of highly active antiretroviral therapy. *Curr Opin Infect Dis* 2006;**19**:26–32.

COMMENTARY

The UK has one of the most extensive HIV registry systems internationally. The study by Dougan and colleagues is an impressive exercise to explore what we can learn from such registries.¹ Still, in the end it remains challenging to conclude whether increased uptake of HIV testing, a rising HIV incidence or both have contributed to the increasing number of HIV diagnoses among (older) men who have sex with men (MSM) using ecological comparisons. A useful addition that may shed more light on this issue would be to construct a mathematical model that incorporates observed data. Such models have been successfully used in the past to predict HIV spread and to assess the influence of strongly interlinked parameters.^{2–4}

The data presented show a stable HIV incidence among MSM in the UK. A rise in HIV incidence is not unexpected considering ongoing high levels of risky sexual behaviour and sexually transmitted infection (STI) epidemics among MSM internationally. To measure the HIV incidence in a population, however, provides a methodological and logistical challenge. The current study uses the serologic testing algorithm for determining recent HIV seroconversion (STARHS) approach that, when incorporated in existing STI/HIV screening programmes at genitourinary medicine (GUM) clinics, is an easy tool to directly estimate HIV incidence. Its wider international (European) application, when standardised, would be of great benefit to HIV incidence surveillance, considering the puzzling discrepant incidence trends found in various countries.

The UK is like, for example, The Netherlands, a country with a historically conservative HIV testing policy. This likely resulted in the still lower testing rates than those found in MSM in, for example, the US or Australia, where testing has been promoted since the beginning of the epidemic.^{5–7} After the introduction of highly active antiretroviral therapy (HAART), the UK and The Netherlands changed to an active approach followed by higher testing uptake. Recently several countries, including the UK, have adopted or are planning to implement the opting-out strategy for HIV testing. This strategy has been shown to drastically reduce the number of undiagnosed HIV infections.^{8–10} Considering that a substantial proportion of patients (one in five MSM in the UK and The Netherlands, and even higher among heterosexuals) are diagnosed late in their infection, the opting-out strategy may also help to diagnose people earlier, when they have a better chance for optimal treatment.^{11–13}

Finally, this study shows the importance of differentiating by age, and agrees with other reports showing an increase in the median age of MSM at HIV diagnosis and showing that HIV incidence is no longer highest in the younger age groups.^{14,15} Although conventionally attention is mainly focused towards the young, older MSM should be specifically targeted in HIV and STI prevention.

Correspondence to: Dr N H T M Dukers, Health Service Amsterdam, Nieuwe Achtergracht, 100 Amsterdam 1018 WT, The Netherlands; ndukers@ggd.amsterdam.nl

REFERENCES

- 1 Dougan S, Elford J, Chadborn T, *et al.* Does the recent increase in HIV diagnoses among men who have sex with men in the United Kingdom reflect a rise in HIV incidence or increased uptake of HIV testing? *Sex Transm Infect* 2007;**83**:120–5.

- 2 **Xiridou M**, Geskus R, de Wit J, *et al*. Primary HIV infection as source of HIV transmission within steady and casual partnerships among men who have sex with men. *AIDS* 2004;**18**:1311–20.
- 3 **Porco TC**, Martin JN, Page-Shafer KA, *et al*. Decline in HIV infectivity following the introduction of highly active antiretroviral therapy. *AIDS* 2004;**18**:81–8.
- 4 **Bezemer DO**, de Wolf F, Boerlijst MC, *et al*. Despite HAART, HIV-1 is once again spreading epidemically amongst men having sex with men in the Netherlands [V-143]. *14th Conference on Retroviruses and Opportunistic Infections (CROI 2007)*. Los Angeles, CA, USA.
- 5 **Stolte IG**, de Wit J BF, Kolader M, *et al*. Low HIV-testing rates among younger high-risk men who have sex with men in Amsterdam. *Sex Transm Infect* 2007. In press.
- 6 **Jin FY**, Prestage G, Law MG, *et al*. Predictors of recent HIV testing in homosexual men in Australia. *HIV Med* 2002;**3**:271–6.
- 7 **Chen SY**, Gibson, Weide D, *et al*. Unprotected anal intercourse between potentially HIV-serodiscordant men who have sex with men, San Francisco. *JAIDS* 2003;**33**:166–70.
- 8 **Weiser SD**, Heisler M, Leiter K, *et al*. Routine HIV testing in Botswana: a population-based study on attitudes, practices, and human rights concerns. *PLoS Med* 2006;**3**:e261.
- 9 **Mulder-Folkerts DK**, van den Hoek JA, van der Bij AK, *et al*. Less refusal to participate in HIV screening among pregnant women in the Amsterdam region since the introduction of standard HIV screening using the opting-out method [in Dutch]. *Ned Tijdschr Geneesk* 2004;**148**:2035–7.
- 10 **Centers for Diseases Control and Prevention**. Advancing HIV prevention: new strategies for a changing epidemic, 2003. *Morb Mortal Wkly Rep* 2003;**52**:329–32.
- 11 **Van Leth F**, Andrews S, Grinsztejn B, *et al*. for the 2NN study group. The effect of baseline CD4 cell count and HIV-1 viral load on the efficacy and safety of nevirapine or efavirenz-based first-line HAART. *AIDS* 2005;**19**:463–71.
- 12 **Siddique MA**, Hartman KE, Dragileva E, *et al*. Low CD4+ T cell nadir is an independent predictor of lower HIV-specific immune responses in chronically HIV-1 infected subjects receiving highly active antiretroviral therapy. *J Infect Dis* 2006;**194**:661–5.
- 13 **Egger M**, May M, Chene G, *et al*. Prognosis of HIV-1-infected patients starting highly active antiretroviral therapy: a collaborative analysis of prospective studies. *Lancet* 2002;**360**:119–29 [Correction appears in: *Lancet* 2002;**360**:1178].
- 14 **Annual report**. HIV Monitoring Foundation [Stichting HIV Monitoring] Amsterdam 2006. http://www.hiv-monitoring.nl/_site1134/images/Jaarverslag2005.pdf (accessed 11 Feb 2007).
- 15 **Dukers HTM**, Fennema SA, van der Snoek EM, *et al*. Trends in HIV incidence and HIV testing behaviour in men who have sex with men: an overview using three incidence sources, The Netherlands, 1984–2005. *AIDS* 2007;**21**:491–9.